

## ESKIMO ARCTIC

*Oeneis alpina* Kurentzov, 1960  
(Nymphalidae)

**Global rank** G3G4 (01Sep1998)

**State rank** S3 (23Jun2006)

### State rank reasons

Not a common species. Range appears to be limited (based on a limited number of occurrences) but is likely incompletely known. Potential threats are not imminent, but include land use practices which degrade moist tundra and bog habitats, as well as effects of climatic warming.

### Taxonomy

Previously known in North America as *Oeneis excubitor*, which Scott (1986) first suggested might be a subspecies of the Siberian *O. alpina*. Expert examination of the type specimens for both reveals they are one species and *O. excubitor* is now considered a straight synonym (Layberry et al. 1998) for *O. alpina*. No subspecies are recognized by these authors.

### General description

A medium-sized brown butterfly; upperside of wings is orange-brown, with basal two-thirds of both wings darker, especially in males; one to three white-centered eyespots present on forewing and two on hindwing, above and below; underside of hindwings striated dark brown and gray (Scott 1986, Layberry et al. 1998). Females have slightly larger wingspan and lighter, more ochreous background color in the basal area of the upperside of wings than males (Troubridge et al. 1982). Similar in appearance to the Brown Arctic (*O. chryxus*), but hindwing eyespots and brown basal two-thirds of upper wings distinguish *O. alpina* (Scott 1986, Layberry et al. 1998). Appearance of eggs, larvae and pupae not reported.

**Length (mm)** 36-47 (wingspan)

### Reproduction

Males congregate on dry tundra hilltops awaiting receptive females (Scott 1986, Layberry et al. 1998). Females move upward to mate, then fly back down to taiga bogs to oviposit eggs on or near host plants (Philip pers. comm.). Flight period is usually biennial (Scott 1986, Layberry et al. 1998); hence, eggs or larvae apparently overwinter twice before pupating in spring/summer.



### Ecology

One flight period, June to mid-July, apparently biennial (mostly in odd years west of Mackenzie River, even years in Mackenzie River and eastward; Scott 1986, Layberry et al. 1998). Species has a strong compulsion to hilltop (i.e. males collect on hilltops to pursue females) and is far more aggressive than any other Nearctic species in the genus *Oeneis* (Troubridge et al. 1982).

### Migration

Dispersal distance unknown.

### Food

Specific host plants for larvae are unknown, but presumed to be a grass feeder (Philip pers. comm.); host plants for *O. chryxus* include grass species *Danthonia*, *Oryzopsis* and *Phalaris* (Scott 1986). Adult foods, if any, are unknown.

### Phenology

Diurnal, flight period is biennial in early summer.

### Habitat

Breeding habitat is wet grassy tundra and taiga bogs, but males congregate on nearby ridges and rocky areas where mating occurs so both features are essential habitat. Usually seen on dry rocky hilltops, desolate scree slopes and windswept ridges (Troubridge et al. 1982, Scott 1986, Layberry et al. 1998).

### Global range

Eastern Siberia (Chukot and Magadan regions) and low arctic of northeastern Alaska, Yukon and western Northwest Territories as far east as Ford Lake (Scott 1986, Lafontaine and Wood 1997, Tuzov et al. 1997, Layberry et al. 1998, Philip pers. comm.). See Layberry et al. (1998) for Canadian range. Extent of the range in Siberia is unknown.

### State range

Species has been found at several sites in Alaska, all of them along the northern foot of the

Brooks Range from near Anaktuvuk Pass to the Dalton Highway (Philip pers. comm.).

**Global abundance**

Population size unknown, but generally an uncommon species with some high local abundance (Layberry et al. 1998).

**State abundance**

Population size unknown. Very common at Oil Spill Hill on the Dalton Highway, and also at the base of Slope Mt., south of Oil Spill Hill (Philip pers. comm.).

**Global trend**

Unknown.

**State trend**

Unknown. Natural fluctuations in population are apparently common (Philip pers. comm.).

**Global protection**

Species afforded no formal protection.

**State protection**

Species afforded no formal protection.

**Global threats**

No immediate threats to populations or habitat. See State threats.

**State threats**

No immediate threats. Land use practices which degrade moist tundra and bog habitats could be potential threats, including road construction; however, even road construction would likely affect only a small portion of the Alaska population. The effects of climate change and warming temperatures in the arctic are unknown, but could include changes to preferred tundra and bog habitats and potential extension of more southerly butterfly species into *O. alpina*'s range.

**Global research needs**

Investigate general life history including host plant species, habitat associations, metamorphic stages and reproductive cycle.

**State research needs**

Investigate host plant species and habitat associations; describe metamorphic stages and reproductive cycle. Study effects of increasing temperatures on tundra and bog habitats.

**State inventory needs**

Determine extent of range in Alaska. Monitor populations biannually during flight periods to identify any trends in distribution or abundance.

**State conservation and management needs**

No imminent threats to this species, so research on life history and inventory/monitoring to determine range and assess population trends should be a priority. Surveys for adults, larvae and eggs could be combined with vegetation surveys in northern Alaska.

**LITERATURE CITED**

- Lafontaine, J.D. and D.M. Wood. 1997. Butterflies and moths (Lepidoptera) of the Yukon. Pp. 723-785 in: Danks, H.V. and J.A. Downes (Eds.). Insects of the Yukon. Biological survey of Canada (terrestrial arthropods), Ottawa, ON.
- Layberry, R.A., P.W. Hall and J.D. Lafontaine. 1998. The butterflies of Canada. University of Toronto Press, Toronto, ON. 280 pp. + color plates.
- Philip, K. 2005. Personal communication via email between Kenelm Philip, University of Alaska, Fairbanks, AK and Jodi McClory, Alaska Natural Heritage Program, Anchorage, AK.
- Scott, J.A. 1986. The butterflies of North America: a natural history and field guide. Stanford University Press, Stanford, CA. 583 pp.
- Troubridge, J.T., K.W. Philip, J.A. Scott and J.H. Shepard. 1982. A new species of *Oeneis* (Satyridae) from the North American Arctic. The Canadian Entomologist 114:881-889.
- Tuzov, V.K., P.V. Bogdanov, A.L. Devyatkin, L.V. Kaabak, V.A. Korolev, V.S. Murzin, D.G. Samodurov, and E.A. Tarasov. 1997. Guide to the butterflies of Russia and adjacent territories (Lepidoptera, Rhopalocera). Vol. 1: Hesperiiidae, Papilionidae, Pieridae, Satyridae. Pensoft Publishers, Sofia – Moscow.

---

## Acknowledgements

### **State Conservation Status, Element Ecology & Life History**

**Author(s):** McClory, J.G. and T.A.  
Gotthardt, Alaska Natural Heritage  
Program, University of Alaska Anchorage, 707 A  
Street, Anchorage, AK,  
<http://aknhp.uaa.alaska.edu>.



**State Conservation Status, Element Ecology  
& Life History Edition Date:** 23Jun2006

**Reviewer(s):** Kenelm Philip, Institute of Arctic  
Biology, University of Alaska Fairbanks,  
Fairbanks, AK.

Life history and global level information were  
obtained from the on-line database, NatureServe  
Explorer ([www.natureserve.org/explorer](http://www.natureserve.org/explorer)). In  
many cases, life history and global information  
were updated for this species account by Alaska  
Natural Heritage Program zoologist, Tracey  
Gotthardt. All global level modifications will be  
sent to NatureServe to update the on-line  
version.

**NatureServe Conservation Status Factors  
Edition Date:** 17Jul1998

**NatureServe Conservation Status Factors  
Author:** Schweitzer, D.F.

**Global Element Ecology & Life History Edition  
Date:** 21May2001

**Copyright Notice:** Copyright © 2005  
NatureServe, 1101 Wilson Boulevard, 15th Floor,  
Arlington Virginia 22209, U.S.A. All Rights  
Reserved. Each document delivered from this  
server or web site may contain other proprietary  
notices and copyright information relating to that  
document.

**Photo credit:** obtained online at:  
[http://www.cbif.gc.ca/spp\\_pages/butterflies/comparaes/ChryxusArcticcomp\\_e.php](http://www.cbif.gc.ca/spp_pages/butterflies/comparaes/ChryxusArcticcomp_e.php).